Available online at www.joac.info

ISSN:

Journal of Applicable Chemistry, 2012, 1 (1): 70-78



(International Peer Reviewed Journal)



Kinetics and Mechanism of the Oxidation of DL-Methionine by Quinolinium Chlorochromate

Ajay Sharma¹, A. Meena¹, J. Khatri¹, P. Swami² and Vinita Sharma^{*1}

Department of Chemistry, J.N.V. University, Jodhpur, 342 005, India.
 Department of Applied Sciences, Global College of Technology, Sitapura, Jaipur, India
 E-mail: drvsharma29@gmail.com

ABSTRACT

The oxidation of methionine (Met) by tetrakis (pyridine) silver dichromate (QCC) in dimethylsulphoxide (DMSO) leads to the formation of corresponding sulphoxide. The reaction is first order with respect to QCC. Michaelis-Menten type kinetics was observed with respect to methionine. The reaction is catalysed by hydrogen ions. The hydrogen-ion dependence has the form: $k_{obs} = a + b \ [H^+]$. The effect of ninteensolvents on oxidation is fitted in Kamlet's and Swain's multiparametric equations. This Study indicates cation-solvating power of the solvent. A suitable mechanism has been postulated.

Keywords: Halochromate, Kinetics, Kamlet's and Swain's model, Methionine, Oxidation.