Available online at www.joac.info



Journal of Applicable Chemistry

2014, 3 (1): 372-378

(International Peer Reviewed Journal)



ISSN: 2278-1862

Hydrotropes as effective reaction Media for the Synthesis of Rafoxanide

Jagannath J. Kadam¹, Vijay Bhave², Gayatri Bharabde³ & Manohar V Lokhande^{4*}

- 1. Department of Chemistry, Bharathi Vidyapeeth College of engineering, NaviMumbai-400614, Maharashtra, INDIA
- Department of Chemistry, Bhavan's College, Andheri (W), Mumbai- 400058, Maharashtra, INDIA
 Department of Chemistry, Institute of Science, fort, Mumbai- 400032, Maharashtra, INDIA
- 4. Department of Chemistry, Sathaye College, Vile Parle (E), Mumbai-400057, Maharashtra, INDIA

Email: manohar2210@gmail.com

Accepted on 20th December 2013

ABSTRACT

The paper describes the synthesis of Rafoxanide. It is used as anthelmintics that expel parasitic worms (Helminthes) from the body or kill them. A traditional remedy of this type is often called a vermifuge. The synthesis of Rafoxanide was carried out using four different hydrotropes. The effect of various hydrotropes on yield, rate constant and activation energy at various temperatures and concentrations are studied.

Keywords: Hydrotropes, Xylene sulfonic acid (XSA), Cumene sulfonic acid (CSA), (n-BBSA): n-Butyl benzene sulfonic acid (n-BBSA), Isobutyl benzene sulfonic acid (I-BBSA), ¹H NMR, IR.