



**Preparation, Characterization study of 2-(2-hydroxy-4-(phenyldiazenyl)benzylidene)-N-(4-phenylthiazol-2-yl) hydrazinecarboxamide and its Metal Complexes**

**Mahadev D Udayagiri<sup>1,2</sup>, Mahendra Raj K<sup>1,3</sup>, Nagesh Gunavanthrao Yernale<sup>1,4</sup>, Shiva Kumar K<sup>2</sup> and B.H.M. Mruthyunjayaswamy<sup>1\*</sup>**

1. Department of Studies and Research in Chemistry, Gulbarga University  
Kalburagi - 585 106, Karnataka, **INDIA**

2. Government College, Sedam Road, Kalburagi - 585 106, Karnataka, **INDIA**

3. BLDE Association's, PG Department of Chemistry, S.B. Arts & K.C.P. Science College,  
Vijayapur - 586 103, Karnataka, **INDIA**

4. G.E. Society's, Aditya Women's Science and Commerce Degree College, Bhalki - 585 328  
Karnataka, **INDIA**


E-mail: [bhmmswamy53@rediffmail.com](mailto:bhmmswamy53@rediffmail.com),

Accepted on 2<sup>nd</sup> March, 2018

**ABSTRACT**

*The Schiff base ligand (L) 2-(2-hydroxy-4-(phenyldiazenyl)benzylidene)-N-(4-phenylthiazol-2-yl)hydrazinecarboxamide obtained by the condensation of N-(4-phenylthiazole-2-yl)hydrazinecarboxamide with 2-hydroxy-4-(phenyldiazenyl)benzaldehyde and its newly synthesized Cu(II), Co(II), Ni(II) and Zn(II) complexes have been characterized by elemental analysis, molar conductance, magnetic susceptibility, thermal analysis and various spectral studies like FT-IR, <sup>1</sup>H NMR, ESI mass, UV-Visible, ESR spectroscopy and powder X-ray diffraction data. The spectral studies confirmed tridentate ONO donor binding of the ligand involving oxygen atom of amide carbonyl, azomethine nitrogen and oxygen of hydroxyl via deprotonation. Spectral analysis indicates octahedral geometry for Cu(II), Co(II) and Ni(II) complexes and tetrahedral geometry for Zn(II) complex. Newly synthesized ligand and its metal complexes were screened for their antibacterial and antifungal activity by minimum inhibitory concentration (MIC) method. The DNA cleavage activities were studied using plasmid DNA pBR322 as a target molecule by agarose gel electrophoresis method. Furthermore, the antioxidant activity of the ligand (L) and its metal complexes were determined in vitro by reduction of 1,1-diphenyl-2-picryl hydrazyl (DPPH), the ligand exhibited more than potent in vitro-antioxidant activity than its metal complexes.*

## Graphical Abstract



Synthesis of 2-(2-hydroxy-4-(phenyldiazenyl)benzylidene)-N-(4-phenylthiazol-2-yl) hydrazinecarboxamide

**Keywords:** Transition Metal complex, Thiazole Schiff base, 2-hydroxy-4-(phenyldiazenyl) benzaldehyde, Biological activity.

---