



Journal of Applicable Chemistry

2014, 3 (6): 2622-2628

(International Peer Reviewed Journal)



Novel Studies of 30M Metal Salts on the Self Condensation of Cyclic Ketones

Barun Kumar Mehta¹, N. Chandra Sekhar², Dandela Rambabu³, Kazumichi Yanagisawa⁴,
Hiyoshizo Kotsuki⁵ and Mandava Venkata Basaveswara Rao^{2*}

1. Department of Chemistry, National Defence Academy, Khadakwasla, Pune-411023, Maharashtra, **INDIA**

2. Department of Chemistry, Krishna University, Machilipatnam-521001, A. P, **INDIA**

3. Department of Chemistry and the National Institute for Biotechnology in the Negev,
Ben-Gurion University of the Negev, Be'er-Sheva 84105, **ISRAEL**

4. Research Laboratory of Hydrothermal Chemistry, Faculty of Science, Kochi University,
Akebono-cho, Kochi 780-8520, **JAPAN**

5. Laboratory of Natural Products Synthesis, Faculty of Science, Kochi University, Akebono-cho,
Kochi 780-8520, **JAPAN**

Email: professormandava@gmail.com, vbrmandava@yahoo.com

Accepted on 24th November 2014

ABSTRACT

A new and effective method was developed for the self-condensation of cyclic ketones in presence of 30M metal salts under hydrothermal conditions.

Keywords: Self-condensation, cyclic ketones, metal salts.

INTRODUCTION

Self-condensation of aldehydes is a well known process[1] but self-condensation of ketones is limited to a few examples[2]. Steric compression accounts for the poor reactivity of ketones. Special reagents and reaction conditions are required for their practical conversion, especially for non-activated cyclic and higher molecular weight ketones [2–7]. Ketones being moderate carbonacids require strong bases, like lithium diisopropylamide and sodium hydride, for their self-condensation[2]. Strong acids, like hydrochloric acid[3a] and polyphosphoric acid[3b] are also known to promote the self-condensation of ketones. Alternative methods reported for the self-aldol condensation of aromatic and aliphatic ketones require organometallic[4,5] or titanium alkoxides [6] while cyclic ketone self-condensation has been reported using a W(CO)₆/CCl₄/UV system [8]. The cationic rhodium complex [Cp*Rh(h₆-C₆H₆)](BF₄)₂ is also known to promote the self-condensation of ketones [7]. While efficient, some of these methods produce a significant quantity of hazardous metals and noxious solvents. The lack of a general strategy for the selective selfcondensation of non-activated ketones under mild conditions limit its application in organic synthesis [2]. Interestingly, during process development for the synthesis of arylidene and alkylidene carbonyl compounds, Arnold *et al.* observed a very slow self-condensation of cyclopentanone by using two molar equivalents of anhydrous lithium perchlorate over a period of 11 days at room temperature [9a]. Recently microwave-assisted solvent-free self-condensation of carbonyl compounds was developed using catalytic amounts of triethylamine and lithium perchlorate [9].

Increasing environmental concern around energy efficiency and wastemanagement provides an opportunity to develop even more powerful and greener methods for well known organic transformations. The hydrothermal reaction is undoubtedly quite effective as a new tool for promoting the desired products. We were particularly interested in the use of aqueous reaction media under hydrothermal conditions [10]. We reported previously the self-condensation of cyclic ketones in hot aqueous ammonium chloride provided novel pyridine derivatives [11]. In the course of our own efforts in this field, now in this paper we describes aqueous metal salts effect on the self-condensation of cyclic ketones under hydrothermal conditions.

MATERIALS AND METHODS

General methods: All reactions requiring anhydrous conditions were conducted in flame-dried apparatus under an inert atmosphere using standard techniques. Syringes and needles for the transfer of reagents were dried at 100°C and allowed to cool in a desiccator over P₂O₅ before use. All the reagents and solvents were reagent grade and were used without further purification unless otherwise specified. Reactions were monitored by thin layer chromatography using aluminum plates pre-coated with a 0.25 mm layer of silica gel containing a fluorescent indicator (Merck Art. 5544); compounds were visualized by UV lamp and ninhydrin spray. Kieselgel 60 (40–63 μm) and neutral aluminum oxide (Acros, 50–200 μm) were used for column chromatography. Chemical shifts (δ) are given in parts per million (ppm). The splitting patterns are reported as s (singlet), d (doublet), t (triplet) and multiplet (m). Coupling constants (*J*) are given in Hz. All reactions were conducted in a Hasteloy-C autoclave reaction vessel with cone and thread fittings and an internal volume of 20 mL, designed to withstand temperatures up to 400°C.

General procedure for Preparation of 2a-2h (entry 1-8): A mixture of ketone (**1a-h**, 4.5 mmol), lithium chloride (0.15 mol), and water (5 mL) was placed in a reaction vessel, and reacted at 250 °C for 24-72 h as shown in table 2. After quenching with water, the mixture was extracted with AcOEt. The crude product was purified by silica gel column chromatography (hexane/AcOEt = 9: 1) to afford **2a-h** in moderate to good yields.


Selected compounds spectral data.

[1,1'-bi(cyclohexylidene)]-2-one (2a): FT IR (neat) (ν , cm⁻¹): 2922, 2851, 1675, 1639, 1444, 729; ¹H NMR (400 MHz, CDCl₃): δ 5.38 (t, *J*=10 Hz, 1H), 2.88 (dd, *J* = 11.2 and 5.0Hz, 1H), 2.42-2.36 (m, 1H), 2.34-2.26 (m, 1H), 2.06-1.96 (m, 4H), 1.96-1.80 (m, 4H), 1.74-1.52 (m, 6H); ¹³C NMR (100 MHz, CDCl₃): δ 211.82, 136.07, 123.88, 58.98, 42.33, 31.05, 27.88, 27.49, 25.50, 25.06, 23.06, 22.62.

2-cyclopentylidenecyclopentan-1-one (2f): FT IR (neat) (ν , cm⁻¹): 2955, 2870, 1706, 1636, 725; ¹H NMR (400 MHz, CDCl₃): δ 2.77 (m, 2H), 2.51 (m, 2H), 2.29 (m, 4H), 1.90 (m, 2H), 1.76-1.64 (m, 4H); ¹³C NMR (100 MHz, CDCl₃): δ 207.57, 158.82, 128.11, 40.02, 34.51, 32.77, 27.17, 26.91, 25.45, 20.31.

RESULTS AND DISCUSSION

Table1. Optimization and metal salts effect for the self condensation of cyclohexanone



Entry	aq MX	Yields (%) ^a 1	Yields (%) ^a 2	Yields (%) ^a 3	Yields (%) ^a 4	Yields (%) ^a 5
1	-----	100	-	-	-	-
2	16MLiCl ^b	37	52	4	6	1
3	30MLiCl	8	77	7	6	2
4	30MNaCl	51	41	5	2	1
5	30MKCl	29	62	4	3	2
6	30MLiBr	21	34	2	38	5
7	30MCaCl ₂	18	52	3	19	8

^a Determined by GC. ^b Saturated solution.

To establish the optimum conditions, we first examined the reaction of cyclohexanone (**1**) without aqueous lithium chloride at 250 °C. No product was obtained. Then we examined the self-aldol condensation reaction of cyclohexanone (**1**) with 16M aqueous lithium chloride which after 24 h gave **2** in an isolated yield of 52% (entry 2). The best results were obtained in the presence of 30 M aqueous lithium chloride, which after 24 h gave **2** in an isolated yield of 77% (entry 3). Other products trimers and tetramers were present only in a very limited amount (about 5–10%, entry 2), as observed in the GC–MS spectra shown in fig 1. It is important to notice that cyclohexanone dimerizes to produce mainly a β,γ -unsaturated ketone. In order to investigate the effect of metal salts, varieties of other metal salts were used and surprisingly, we observed effect on the yield or ratios of the products (**2-5**) were different (shown in table 1, entry, 4-7). The reaction didn't proceed when the use of 2-methylcyclohexanone instead of cyclohexanone (scheme-1).






Fig. 1. GCMS Data (Column: ULBON HR-54)














Scheme 1. Self condensation of cyclic ketones.

By using optimised condition we used different cyclic ketones **1a-h** (entry 1-8) to produce self condensation products **2a-h** (entry 1-8) in moderate to good yields as shown in table 2. It is important to notice that cyclopentanone (**1f**, entry **6**, table 2) dimerizes to yield predominantly an α,β -unsaturated ketone (**2f**, entry **6**, table 2), while cyclohexanone (**1a**, entry **1**, table 2) dimerizes to produce mainly a β,γ -unsaturated ketone (**2a**, entry **1**, table 2). The difference in the geometry of these two starting materials may be the key factor behind the observed difference in reactivity. Due to the thick nature and high boiling liquid, cycloheptanone and cyclooctanone gave low yield of the corresponding dimmers (**2g** & **2h**, entries **7** & **8**, table 2). Reaction mechanism for the formation of products **2** and **3** shown in scheme 2.

Table2. Substrate Scope

Entry	Ketone	Conditions	Products (major)	Yields (%) ^a
1	 1a	250 °C, 24 h	 2a	77
2	 1b	250 °C, 72 h	 2b	56
3	 1c	250 °C, 72 h	 2c	25
4	 1d	250 °C, 24 h	 2d	36

5	 1e	250 °C, 72 h	 2e	57
6	 1f	250 °C, 24 h	 2f	38
7	 1g	250 °C, 48 h	 2g	29
8	 1h	250 °C, 72 h	 2h	21

^b Isolated yields.

Scheme 2. Probable reaction mechanism leading to product 2 and 3.

APPLICATIONS

The obtained results illustrate the potential utility of this method as an environmentally-friendly process.

CONCLUSIONS

In summary, we have developed a novel and effective method for the self-condensation of cyclic ketones using aqueous metal salts under hydrothermal conditions. The present methodology has described a remarkable metal salts effect on the self-condensation of cyclic ketones and results illustrate the potential utility of this method as an environmentally-friendly process. Further studies to explore the possibility of hydrothermal reactions in other types of transformation are in progress.

Conflict of Interest: The authors confirm that this article content has no conflict of interest.

ACKNOWLEDGEMENTS

The authors thank University Grants Commission for the financial assistance under Major Research Project.

REFERENCES

- [1] Y.Watanabe, K. Sawada, M. Hayashi, A green method for the self-aldol condensation of aldehydes using lysine, *Green Chem.*, **2010**, 12, 384-386.
- [2] (a) H.O. House, D.S. Crumrine, A.Y. Teranishi, H.D. Olmstead, Chemistry of carbanions, XXIII. Use of metal complexes to control the aldol condensation, *J. Am. Chem. Soc.*, **1973**, 95(10), 3310-3324; (b) S.E. Denmark, W. Lee, Investigations on transition state geometry of the aldol condensation in aqueous medium, *Tetrahedron Lett.*, **1992**, 33(50), 7729-7732; (c) F. Fringuelli, G. Pani, O.Piermatti, F. Pizzo, Condensation-reactions in water of active methylene compounds with arylaldehydes one pot synthesis of flavonols, *Tetrahedron*, **1994**, 39, 11499-11508; (d) R. Mestres, A green look at the aldol reaction, *Green Chem.*, **2004**, 6, 583-603; (e) C.Capello, U. Fischer, K. Hungerbuhler, What is a green solvent? A comprehensive framework for the environmental assessment of solvents, *Green Chem.*, **2007**, 9, 927-934.
- [3] (a) A.T. Nielsen, W. Houlihan, *J. Org. React.*, John Wiley, NewYork, **1968**, vol 16, p. 1; (b) A.R. Bader, US Patent no. 2769842,1956; A. R. Bader, *Chem. Abstr.*, **1956**, 52, 439.
- [4] T. Nakano, S. Irefune, S. Umamo, A. Inada, Y. Ishii, M. Ogawa, Ruthenium-catalyzed selective addition of carboxylic acids to alkynes, A novel synthesis of enol esters, *J. Org. Chem.*, **1987**, 85(11), 2239-2244.
- [5] C.Danen, T.T. Kensler, The Condensation of Ketones by Aluminum *t*-Butoxide to Compounds of the Mesityl Oxide Type, *J. Am. Chem. Soc.*, **1940**, 62(12), 3401-3404.
- [6] Y.G.Yatluk, V.Y.Sosnovskikh, A.L. Suvorov, Condensation of Ketones in the Presence of Titanium Alkoxides, *Russ. J. Org. Chem.*, **2004**, 40(6), 763-765.
- [7] H. Terai, H. Takaya, T. Naota, Rhodium-catalyzed direct aldol condensation of ketones: a facile synthesis of fused aromatic compounds, *Tetrahedron Lett.*, **2006**, 47(11), 1705-1708.
- [8] C.Bozkurt, Aldol-type condensation reactions of cyclic ketones by the W(CO)₆/CCl₄/UV system, *J. Organomet. Chem.*, **2000**, 603, 252.
- [9] L.K.Sharma, K.B. Kimb, G.I. Elliott. A selective solvent-free self-condensation of carbonyl compounds utilizing microwave irradiation, *Green Chem.*, 2011, **13**, 1546-1549.
- [10] (a) B.K. Mehta, S. Dhar, H. Ila, H.Junjappa, Highly Efficient and Selective Displacement of Alkylthio Group on Acylketene O, S-Acetals by Organocopper Reagents: A Novel Route to 2-Alkoxy/Aryloxy-1-Alkenylketones, *Tetrahedron Lett.* **1995**, 36, 9377-9380. (b) B.K.Mehta, K. Yanagisawa, M.Shiro, H. Kotsuki, Unprecedented Hydrothermal Reaction of O-Phenylaniline and

- Related Derivatives with Cyclic Ketones. A Novel Approach to the Construction of Phenanthridine and Quinoline Ring Systems, *Organic Lett*, **2003**, 5, 1605-1608.
- [11] H. Kotsuki, B. K. Mehta, K. Yanagisawa, Novel Synthesis of Fully-Substituted Pyridine Derivatives via Self-Condensation of Cyclic Ketones in Aqueous Ammonium Chloride under Hydrothermal Conditions, *Synlett* **2001**, 8, 1323–1325.