



## Synthesis and Characterization of Iron (III) Complex Containing Schiff Bases Derived from Sulfadiazine

Ahmed Neamah Thamer Al-Yasiry

The General Directorate of Educational in Najaf Al-Ashraf,  
Dr. Enad Ghazwan Preparatory School for Boys, **IRAQ**

Email: [ahmedalyasiry@yahoo.com](mailto:ahmedalyasiry@yahoo.com)

Accepted on 7<sup>th</sup> November 2017, Published online on 27<sup>th</sup> November 2017


### ABSTRACT

Schiff base compound [(4-((2-hydroxynaphthalen-1-yl) methylene amino)-N-(pyrimidin-2-yl) benzene sulfonamide) (L) was synthesized by reaction Sulfadiazine with 2-hydroxy naphthaldehyde in glacial acetic acid as solvent and catalyst. New complex (Lx) of Fe (III) with organic ligand Schiff base was also prepared. Ligand and complex were identified by elemental analysis, magnetic moment, <sup>1</sup>H-NMR spectrum and molar conductance along with electronic and infrared spectral analysis. Octahedral geometry around this metal ion has been proposed on the basis of magnetic and spectral studies.

### High Lights:

1. Synthesis and characterization new ligand [(4-((2-hydroxynaphthalen-1-yl)methyleneamino)-N-(pyrimidin-2-yl)benzenesulfonamide]
2. Synthesis New complex of Fe(III)with organic ligand Schiff base
3. Study magnetic and spectral properties of ligand and complex
4. Study anti-bacterial activity of ligand and complex

### Graphical Abstract:



**Keywords:** Sulfadiazine, Schiff base, Ligand and complex.


## INTRODUCTION

Sulfadiazine [4-amino-N-2-pyrimidinylbenzene sulfonamide [1] is one of sulfonamide group of antibiotics that using as a veterinary medicine [2]. It eliminates type of bacteria which cause infections by preventing preparation of folic acid which bacteria was needed for growth. It is used for the treatment of many of infections for example toxoplasmosis, malaria, urinary tract infections and a certain type of brain infections. Sulfa drugs were possessing (-SO<sub>2</sub>NH-) Moiety which is very important toxophoric functional group' concede chemotherapeutically important such as sulfadiazine (silver sulfadiazine), sulfathiazole, sulfamerazine [3]. Generally, there are two reactive positions in sulfadiazine, the first sulfonamide group and the second aromatic amine [4]. Schiff bases are the organic compounds containing azomethine group (HC=N-) synthesized by condensation of a primary amine with (aldehyde or ketone) an carbonyl compound [5]. They were first reported by Schiff in 1864 [6] and the formula of the Schiff base is (RR'C=NR'') where R, R', R'' is an alkyl group or phenyl group that makes Schiff bases a stable imine [7]. Schiff bases are the important organic compounds showed wide spectrum of biological activities and many applications [8]. The common oxidation states shown by iron are +2 and +3. Almost all the aqueous chemistry of iron is confined to the +2 and +3 oxidation states. Fe (III) ions form a large number of complexes than Fe (II) ion. In most of these, the coordination number of Fe (III) ion is six and the complexes formed are octahedral [9]. Thus, Schiff base has good chelating ability with metal ions and show various absorption bands differ in position as well as intensity [10,11]. In view of these factors, in this research synthesized new complex of Fe (III) ions with linked Schiff bases containing oxygen and nitrogen donor atoms.

## MATERIALS AND METHODS

All chemicals and solvents were highest purity obtained from Fluka. The Merck and BDH. The melting points were determined on a Electrothermal, melting point 9300. Elemental analysis (C.H.N) was obtained using Euro VECTOR (C.H.N) elemental analyzer. IR spectra were recorded using KBr discs in the range (4000-400) cm<sup>-1</sup> on FTIR Test scan Shimadzu model 8300, while the (UV-Vis). Spectra recorded in ethanol on Shimadzu model 1650PC. Molar conductance measurements were carried out at lab temperature in DMF (10<sup>-3</sup> M), HANNA model 214EC conductivity meter. The pH measurements by using pH-meter Hanna. The metal content of the complex was determined by using atomic absorption technique by Perkin-Elmer model 2280.


**Preparation of ligands:** (4mmol, 1g) Sulfadiazine dissolved in (20 mL) glacial acetic acid and 2-hydroxy-1-naphthaldehyde (4mmol, 0.688 g) was added drop wise. The resulting mixture was stirred at (80<sup>o</sup>C) for (14) h. Then, the TLC showed that the reaction was completed by using (hexane: ethylacetate, 4:1). The solution was separated by using diethyl ether. The organic layers were dried over MgSO<sub>4</sub>. Removal of the solvent and recrystallize. The characteristics are - (4-((2-hydroxynaphthalen-1-yl)methyleneamino)-N-(pyrimidin-2-yl)benzenesulfonamide is liquid, Chemical Formula: C<sub>21</sub>H<sub>16</sub>N<sub>4</sub>O<sub>3</sub>S; m.p. 240-242, yield 60%; Elemental Analysis: H, 3.99; S, 7.93; C, 62.36; N, 13.85(calcd.); H, 3.89; S, 7.873; C, 62.144; N, 13.778(found); IR (KBr disc, cm<sup>-1</sup>): 3340 (N-H sulfonamide Str), 3070 (Aryl C-H), 2944, 2843 (Alkyl C-H), 1661 (C=O str), 1537 (C=N str), 1312 (SO<sub>2</sub>sulfonamide sym.) 1147 (SO<sub>2</sub>Asym.str.), 1425-1600 (Aromatic ring); <sup>1</sup>HNMR (δ ppm), (DMSO-*d*6) : 11.321 (SO<sub>2</sub>-NH), 8.724 (HC=N pyrimidine ring), 4.003 (C-H triazoline ring), 8.615-6.625 ( Aryl C-H), 3.19 (N-(CH<sub>3</sub>)<sub>2</sub>).



Scheme -1

**Preparation of metal complex:** To a hot solution of ligand (2 mmol) in 30mL methanol, the ferric chloride (1 mmol) in methanol was added drop wise. The pH of the solution was adjusted to 7 to 8.5 by adding alcoholic ammonia. The contents were digested for 4 h. The Fe (III) complexes were filtered in hot condition, product washed several times with hot methanol and then by petroleum ether and dried in vacuum. Chemical Formula:  $[\text{Fe}(\text{C}_{42}\text{H}_{30}\text{Cl}_2\text{N}_8\text{O}_6\text{S}_2)]$ ; Elemental Analysis: C, 54.03; H, 3.24; N, 12.00; S, 6.87; Fe, 5.98(calcd.), C, 53.889; H, 3.109; N, 11.977; S, 6.721; Fe, 5.78(found).

The structure of metal complexes is shown below.



Structure of Fe(III)Complex with Ligand

## RESULTS AND DISCUSSION

Ligand derived from reaction of Sulfadiazine with 2-hydroxy naphthaldehyde in glacial acetic acid. The [C.H.N.S] analysis of synthesized ligand was accepted agreement with the calculated percentage of elements.  $^1\text{H-NMR}$  spectrum considers good evidence for prepared ligand (Figure 1). The complex is colored solid and stable to air. It is insoluble in methanol, ethanol and water but soluble in DMF, DMSO. The synthesized ligand (L) was characterized by ultra-violet-visible (UV-Vis) spectroscopic technique and recorded in ethanol on Shimadzu model 1650P. The spectrum of the ligand in ethanolic solution ( $10^{-3}$  M) gave mainly three peaks, at (323 nm) and (377 nm) due to the moderate energy  $\pi-\pi^*$  transition of the phenyl rings. At the (487 nm) peak ( $\lambda_{\text{max}}$ ) due to the  $\pi-\pi^*$  transition which take place from benzene ring by intermolecular charge-transfer through the azomethine group (Schiff base) ( $-\text{C}=\text{N}$ ). Interaction of the (Fe III) ion with the ligand (L) has been studied in solution; An aqueous ethanolic solutions were always performed over wide molar concentration and acidity range. The interaction of the iron ion (Fe metal) with the ligand manifest itself in the absorption spectra by the appearance of a peak in the range (511 nm). A great bathochromic shift obtained in the visible region was determined in the complex solutions spectra with respect to that of the ligand alone. The high shift in the ( $\lambda_{\text{max}}$ ) gave a good indication for complex formation. The Infrared spectra of the metal complexes were recorded between  $4000-400\text{ cm}^{-1}$ . In IR spectra of free ligands, the band observed in the range  $1622-1562\text{ cm}^{-1}$  were assigned to the  $\text{C}=\text{N}$  stretching frequencies,  $3326\text{ cm}^{-1}$  (N-H) stretching of sulfonamide,  $3410\text{ cm}^{-1}$  (OH) stretching,  $3070\text{ cm}^{-1}$  (Aryl C-H) stretching,  $1340\text{ cm}^{-1}$  ( $\text{SO}_2$  sulfonamide sym.),  $1122\text{ cm}^{-1}$  ( $\text{SO}_2$  Asym. Stretching). The good

evidence of bonding is revealed by the appearance of band at 477-530  $\text{cm}^{-1}$ (Fe-O), 432- 478  $\text{cm}^{-1}$ (Fe-N) [12]. The molar conductance values of all the complex in DMF at a concentration of  $1 \times 10^{-4}$  M fall in the range 25.44  $\text{mho.cm}^2\text{mol}^{-1}$ . The conductance value is low molar indicates non-electrolytic nature of complex [13]. The Fe (III) complex show magnetic moment values are in the range 3, 22 B.M. at room temperature suggesting high spin octahedral geometry [14].




Fig 1.  $^1\text{H}$ NMR Spectrum of ligand

### APPLICATIONS

Schiff bases (L) and their Fe (III) complex (Lx) was screened (at 300 and 600 ppm) in vitro for their antibacterial activity against of types of bacteria such as Sta. aureus, E. coli and Pse. aeruginosa. The ligand showed 30-60 % inhibition (Table 1). The complex had greater antibacterial activity than the corresponding ligand 40-80 % inhibition.

Table 1. Antibacterial activities of compounds (L and LX)

Comp. No.	Escherichia coli		Staphylococcus aureus		Pseudomonas aeruginosa	
	300 ppm Zone of inhibition (mm)	600 ppm Zone of inhibition (mm)	300 ppm Zone of inhibition (mm)	600 ppm Zone of inhibition (mm)	300 ppm Zone of inhibition (mm)	600 ppm Zone of inhibition (mm)
<b>L</b>	30	50	40	60	35	60
<b>LX</b>	40	70	45	70	50	80

### CONCLUSIONS

In the present study, reported the synthesis of Fe(III) complex of azomethine (Schiff base) ligand derived from reaction of Sulfadiazine with 2-hydroxy naphthaldehyde in glacial acetic acid. Infrared spectral studies suggest the metal-ligand coordination at azomethine and OH group. Further, by elemental analysis, molar conductivity measurements, magnetic susceptibilities, electronic absorption, structure for these Fe(III) complex is assigned as high spin octahedral structure.

## REFERENCES

- [1] M. John; "Organic medicinal and pharmaceutical chemistry", Ed 2, **2004**, p 237.
- [2] D. Miller, R.M. Rashid, L. Falzon, E.M. Elamin and S.Zehtadchi, A definitive case of metastatic folliculotropic melanoma, *J. Am. Acad. Dermatol.*, **2012**, 66(5), 159-165.
- [3] P. A. Ajibade, A.G. Kolawole, P. Obrien, M. Helliwell and J.Raftery; Cobalt (II) complexes of the antibiotic sulfadiazine, the X-ray single crystal structure of  $[\text{Co}(\text{C}_{10}\text{H}_9 \text{N}_4\text{O}_2\text{S})_2 (\text{CH}_3\text{OH})_2]$ , *Inorg. Chim. Acta*, **2006**, 359, 3111–3116.
- [4] Z. Lin, Z. Huang, L. Shi and J. Huang, Synthesis of a new monomer N'-( $\beta$ -methacryloyloxyethyl)-2-pyrimidyl-(pbenzyloxycarbonyl)aminobenzenesulfonamide and its copolymerization with N-phenyl maleimide, *Journal of polymer Science*, **2000**, 38, 2548-2554.
- [5] P. M. Henry and G. L. Lange, "The Chemistry of double bonded functional groups", S. Patai, Interscience, New York, **1977**, 1067.
- [6] Y.Li, Z. S. Yang, H. Zhang, B. J. Cao and F. D. Wang, Artemisinin Derivatives Bearing Mannich Base Group Synthesis and Antimalarial Activity, *Bioorg. and Med. Chem.*, **2003**, 11, 4363-4368.
- [7] C.S. Naveena, B. Poojary, V. Kumar, S. Shenoy, Synthesis, Characterization and Antimicrobial Activities of Some Novel Thiazole-[3,2-b]-1,2,4-triazole Derivatves, *J. Applicable. Chem.*, **2015**, 4, 828-839.
- [8] K. Singh and Ritu, Structural Elucidation of Bio-Active Metal Complexes of Schiff Base Derived from Condensation of 2, 5-Dimethoxybenzaldehyde with Substituted 1, 2, 4-Triazine Scaffold, *J. Applicable. Chem.*, **2017**, 6, 299-310
- [9] B. Kalluraya, K. Sushma, A. Kumar, K. Kaushik, B.R. Aparna and V. Shetty, Synthesis, Characterization and Biological Activities of 1, 2, 3-Triazole Containing Substituted Cyclo hexenones, *J. Applicable. Chem.*, **2017**, 6, 08-18.
- [10] F.A. Cotton and G.Wilkinson, "Advanced Inorganic Chemistry," 6th edition, wiley Eastern, New Delhi, **1999**, P 89-91.
- [11] N.K. Sarmah, K.N. Sarmah, T.V. Patel, Synthesis and Antimicrobial Evaluation of a Novel Series of Some S-Triazine Moiety, *J. Applicable. Chem.*, **2014**, 3, 1936-1944.
- [12] M.B. Halli, S. Suryakant, M. Kinni, Synthesis, Characterization and Biological Activities of Heterocyclic Schiff Base and its Metal Complexes, *J. Applicable. Chem.*, **2015**, 4, 467-475.
- [13] M.R. Anil, S. Shanmukhappa, B.E. Rangaswamy and M. Revanasiddappa, Synthesis, Characterization and Antimicrobial Activity of Some Transition Metal Complexes with Schiff Base Ligand, *J. Applicable Chem*, **2014**, 3(4), 1440-1446.
- [14] A.Ramachandra, P.Nirdosh and S. D. Angadi, Synthesis, Characterization and Microbial Activities of Metal Complexes with Coumarine Derivatives, *Rasayan J. Chem*, **2009**, 2, 1, 81-86.

## AUTHOR ADDRESS

### 1. Ahmed NeamahThamer Al-Yasiry

Assistant Lecturer, The General Directorate of Educational in  
Dr. Enad Ghazwan Preparatory School for Boys  
Najaf Al-Ashraf- Iraq  
E-mail: ahmedalyasiry@yahoo.com, Mobile No 0780308286