



Mechanism of the Oxidation of some α -Hydroxy acids by Imidazolium Dichromate

Archana Sharma¹, Jigyasa Sharma¹, Ruchi Kalla¹,
Gifty Sharma¹, Laszlo Kotai² and Pradeep K. Sharma^{1*}


1. Department of Chemistry J.N.V. University, Jodhpur, Rajasthan, **INDIA**
2. Institute of Material and Environmental Chemistry, RCNS, HAS, Budapest, **HUNGARY**
Email: drpkvs27@yahoo.com

Accepted on 7th November, 2018

ABSTRACT

The oxidation of glycolic, lactic, malic and a few substituted mandelic acids by imidazolium dichromate (IDC) in dimethylsulphoxide (DMSO) leads to the formation of corresponding oxoacids. The reaction is first order each in IDC. Michaelis-Menten type of kinetics is observed with respect to the hydroxy acids. Reaction is failed to induce the polymerization of acrylonitrile. The oxidation of α -deuteriomandelic acid shows the presence of a primary kinetic isotope effect ($k_H/k_D = 5.77$ at 298 K). The reaction does not exhibit the solvent isotope effect. The reaction is catalyzed by the hydrogen ions. The hydrogen ion dependence has the form: $k_{obs} = a + b [H^+]$. Oxidation of p-methyl mandelic acid has been studied in 19 different organic solvents. The solvent effect has been analyzed by using Kamlet's and Swain's multiparametric equations. A mechanism involving a hydride ion transfer via a chromate ester is proposed.

Graphical Abstract



Scheme

Keywords: Correlation analysis, Dichromate, Hydroxy acids, Kinetics, Mechanism, Oxidation.

INTRODUCTION

It has been reported that various salts of Cr (VI) are well known oxidizing reagents in synthetic organic chemistry, but are sometimes non-selective due to their drastic nature as an oxidant. Further, these salts are insoluble in most of the organic solvents. Therefore, solubility is one of the main problems. To avoid various limitations with these compounds, a large number of organic derivatives of Cr (VI) have been prepared and used in organic synthesis as mild and selective oxidants in non-aqueous solvents [1-4]. One of such compounds is imidazolium dichromate (IDC) [5]. We have been interested in the kinetic and mechanistic aspects of the oxidation by complex salts of Cr (VI) and several reports on halochromates and dichromates are available in the literature [6-10]. Therefore, it was of interest to investigate the kinetics of the oxidation of some α -hydroxy acids by IDC in DMSO. A suitable mechanism has also been postulated.

MATERIALS AND METHODS

Materials: The hydroxy acids were commercial products of the highest purity available and were used as such. The preparation and specification of the substituted mandelic acids have been described earlier [11]. IDC was prepared by reported method [5] and its purity was checked by an iodometric method. α -Deuteriomandelic acid (PhCD (OH)COOH or DMA) was prepared by the method of Kemp and Water [12]. Its isotopic purity, ascertained by NMR spectra, was $95 \pm 4\%$. Due the non-aqueous nature of the solvent, toluene p-sulphonic acid (TsOH) was used as a source of hydrogen ions. Solvents were purified by their usual methods [13].

Product analysis: Product analyses were carried out under kinetic conditions i.e., with an excess of the reductant over IDC. In a typical experiment mandelic acid (7.6 g, 0.05 mol) and IDC (3.54 g, 0.01 mol) were dissolved in 100 ml of DMSO and was allowed to stand in dark for ≈ 24 h to ensure the completion of the reaction. It was then treated with an excess (250 mL) of a freshly prepared saturated solution of 2,4-dinitrophenylhydrazine in 2 mol dm^{-3} HCl and kept overnight in a refrigerator. The precipitated 2,4-dinitrophenyl-hydrozone (DNP) was filtered off, dried, weighed, recrystallized from ethanol and weighed again. The product was identical (mp and mixed mp) to an authentic sample of DNP of phenylglyoxylic acid. Similar experiments with the other hydroxy acids yielded the DNP of the corresponding oxoacids in 78 to 87% yields, after Recrystallisation. The oxidation state of chromium in completely reduced reaction mixtures, determined by an iodometric method, was 3.95 ± 0.10 .


Kinetic measurements: The pseudo-first order conditions were attained by keeping a large excess ($\times 15$ or greater) of the hydroxy acid over IDC. The temperature was kept constant to ± 0.1 K. The solvent was DMSO, unless specified otherwise. The reactions were followed by monitoring the decrease in the concentration of IDC spectrophotometrically at 354 nm for up to 80% of the reaction. No other reactant or product has any significant absorption at this wavelength. The pseudo-first order rate constants, k_{obs} , were computed from the linear least square plots of $\log [\text{IDC}]$ versus time. Duplicate kinetic runs showed that the rates were reproducible within $\pm 3\%$. All experiments, other than those for studying the effect of hydrogen ions, were carried out in the absence of TsOH.

RESULTS AND DISCUSSION


The rate and other experimental data were obtained for all the hydroxy acids studied. Since the results were similar, only representative data are reproduced here.

Stoichiometry: The oxidation of hydroxy acids resulted in the formation of the corresponding oxoacids. Product analysis and stoichiometric determinations indicated that the overall reaction could

be written as below (1).



Rate Laws: The reactions are of first order with respect to IDC. Further, the pseudo-first order rate constant, k_{obs} is independent of the initial concentration of IDC. The reaction rate increases with increase in the concentration of the hydroxy acid but not linearly (Table 1). A plot of $1/k_{\text{obs}}$ against $1/[\text{Hydroxy acid}]$ is linear ($r > 0.995$) with an intercept on the rate-ordinate (Figure 1). Thus, Michaelis-Menten type kinetics are observed with respect to the hydroxy acid. This leads to the postulation of following overall mechanism (2) and (3) and rate law (4).



$$\text{Rate} = k_2 K [\text{HA}] [\text{IDC}] / (1 + K [\text{HA}]) \quad (4)$$

Table 1. Rate constants for the oxidation of mandelic acid by IDC at 298 K

$10^3 [\text{IDC}]$ mol dm^{-3}	$[\text{HA}]$ mol dm^{-3}	$10^4 k_{\text{obs}}$ s^{-1}
1.00	0.10	10.6
1.00	0.20	15.7
1.00	0.40	20.5
1.00	0.60	22.9
1.00	0.80	24.3
1.00	1.00	25.2
1.00	1.50	26.5
1.00	3.00	28.0
2.00	0.20	14.4
4.00	0.20	16.2
6.00	0.20	15.3
8.00	0.20	15.0
1.00	0.40	21.6*

*contained 0.001 M acrylonitrile




Figure 1. Oxidation of Mandelic Acid by IDC at 308K: A typical Kinetic Run

The dependence of reaction rate on the reductant concentration was studied at different temperatures and the values of K and k_2 were evaluated from the double reciprocal plots. The

thermodynamic parameters of the complex formation and activation parameters of the decomposition of the complexes were calculated from the values of K and k_2 respectively at different temperatures (Tables 2 and 3). Figure 2 depict a typical kinetic run.




Figure 2. $1/k_{\text{obs}}$ vs $1/[\text{HA}]$: A double reciprocal plot.

Table 2. Formation constants for the decomposition of IDC-hydroxy acid complexes and thermodynamic parameters

R	$K / (\text{dm}^3 \text{mol}^{-1})$				$-\Delta H^*$	$-\Delta S^*$	$-\Delta G^*$
	288 K	298 K	308 K	318 K	(kJ mol^{-1})	($\text{J mol}^{-1}\text{K}^{-1}$)	(kJ mol^{-1})
H	7.21	6.42	5.58	4.77	12.9±0.5	20±2	7.06±0.4
p-F	6.85	6.03	5.25	4.41	13.6±0.5	23±2	6.92±0.4
p-Cl	6.03	5.24	4.42	3.60	15.5±0.7	31±2	6.55±0.6
p-Br	5.85	5.02	4.25	3.42	16.0±.7	32±2	6.45±0.6
p-Me	6.01	5.22	4.37	3.60	15.5±0.6	31±2	6.54±0.5
p-Pr ⁱ	5.94	5.15	4.32	3.53	15.7±0.7	31±2	6.50±0.5
p-OMe	5.90	5.07	4.27	3.49	15.8±0.6	32±1	6.48±0.5
m-Cl	6.30	5.51	4.68	3.88	14.8±0.6	28±1	6.68±0.5
m-NO ₂	5.67	4.85	4.03	3.24	16.6±0.7	35±2	6.36±0.6
p-NO ₂	5.58	4.75	3.97	3.15	16.9±0.8	36±2	6.31±0.6
GA	6.15	5.36	4.50	3.72	15.3±0.6	30±2	6.60±0.5
LA	7.29	6.45	5.67	4.85	12.8±0.4	20±1	7.09±0.3
MLA	6.30	5.50	4.70	3.87	14.8±0.6	28±2	6.68±0.5
DMA	5.99	5.15	4.41	3.60	15.3±0.6	30±2	6.53±0.5

Table 3. Rate constants and activation parameters for the oxidation of hydroxy acid by IDC

R	$10^4 k_2 / (\text{dm}^3 \text{mol}^{-1} \text{s}^{-1})$				ΔH^*	$-\Delta S^*$	ΔG^*
	288 K	298 K	308 K	318 K	(kJ mol^{-1})	($\text{J mol}^{-1}\text{K}^{-1}$)	(kJ mol^{-1})
H	8.55	16.2	29.7	53.1	43.8±0.2	152±1	88.9±0.2
p-F	12.6	22.5	41.4	72.9	42.2±0.6	154±2	88.1±0.5
p-Cl	4.86	9.36	18.0	31.5	45.2±0.4	152±1	90.3±0.3
p-Br	3.96	7.65	14.4	26.1	45.4±0.2	153±1	90.8±0.1
p-Me	42.3	75.6	126	216	38.6±0.4	157±1	85.1±0.3
p-Pr ⁱ	36.0	65.7	117	189	39.8±0.5	154±2	85.5±0.4
p-OMe	486	783	1220	1860	31.5±0.1	161±1	79.3±0.1
m-Cl	1.26	2.52	5.04	9.81	49.6±0.5	148±2	93.5±0.4
m-NO ₂	0.21	0.47	1.04	2.07	55.6±0.1	142±1	97.7±0.1
p-NO ₂	0.15	0.34	0.72	1.53	56.2±0.3	142±1	98.5±0.3
GA	3.78	6.75	12.6	22.5	42.9±0.7	162±2	91.0±0.6
LA	5.40	10.4	20.7	38.7	47.7±0.6	142±2	90.0±0.5
MLA	4.77	9.27	18.0	34.2	47.5±0.5	144±2	90.3±0.4
DMA	1.39	2.81	5.50	10.5	48.8±0.3	150±1	93.3±0.2
$k_{\text{H}}/k_{\text{D}}$	6.15	5.77	5.40	5.06			

Induced polymerization of acrylonitrile: The oxidation of hydroxy acids, by IDC, in an atmosphere of nitrogen failed to induce the polymerization of acrylonitrile. Further, addition of acrylonitrile had no effect on the rate (Table 1). We further confirm the absence of free radicals in the reaction pathway, the reaction was carried out in the presence of 0.05 mol dm⁻³ of 2,6-di-*t*-butyl-4-methylphenol (butylated hydroxytoluene or BHT). It was observed that BHT was recovered unchanged, almost quantitatively.

Effect of acidity: Reaction is catalyzed by hydrogen ions. *p*-Toluene sulphonic acid (TsOH) was used as a source of hydrogen ions. The hydrogen ion dependence has the following form (Table 4). The values of *a* and *b*, for mandelic acid, are 9.91±0.29×10⁻⁴ s⁻¹ and 17.8±0.47×10⁻⁴ mol⁻¹ dm³ s⁻¹ respectively (*r*² = 0.9972).

$$k_{\text{obs}} = a + b [\text{H}^+] \quad (5)$$

Table 4. Effect of hydrogen ion concentration on the oxidation of mandelic acid by IDC

[IDC] = 0.001 mol dm ⁻³ ;	[HA] = 1.0 mol dm ⁻³ ;					Temp. = 298 K
[H ⁺]/mol dm ⁻³	0.10	0.20	0.40	0.60	0.80	1.00
10 ⁴ k _{obs} /s ⁻¹	11.7	13.5	17.4	20.0	24.3	27.9

Kinetic isotope effect: To ascertain the importance of the cleavage of the C-H bond in the rate-determining step, the oxidation of α-deuteriomandelic acid (DMA) was studied. Results showed the presence of a substantial primary kinetic isotope effect (Table 3). The value of *k*_H/*k*_D is 5.77 at 298 K.

Effect of solvents: The rate of oxidation of mandelic acid was determined in nineteen different organic solvents. The choice of the solvents was limited by the solubility of IDC and reaction with primary and secondary alcohols. There was no noticeable reaction with the solvents chosen. The kinetics was similar in all the solvents. The values of *k*₂ at 298 K are recorded in table 5.

Table 5. Effect of solvents on the oxidation of *p*-methyl mandelic acid by IDC at 308 K

Solvents	K (dm ⁻³ mol ⁻¹)	10 ⁴ k _{obs} (s ⁻¹)	Solvents	K (dm ⁻³ mol ⁻¹)	10 ⁴ k _{obs} (s ⁻¹)
Chloroform	6.30	41.7	Toluene	4.90	14.1
1,2-Dichloroethane	5.58	51.3	Acetophenone	4.88	61.7
Dichloromethane	6.21	46.8	THF	5.25	20.4
DMSO	4.37	126	<i>t</i> -Butylalcohol	5.77	15.1
Acetone	4.66	37.2	1,4-Dioxane	4.45	25.1
DMF	5.27	63.1	1,2-Dimethoxyethane	5.78	12.0
Butanone	5.05	33.9	CS ₂	4.68	6.46
Nitrobenzene	5.88	49.0	Acetic Acid	4.96	7.94
Benzene	6.11	18.2	Ethyl Acetate	5.70	15.8
Cyclohexane	6.03	1.86			

The entropy and enthalpy of activation of the oxidation of ten substituted mandelic acids are linearly related (*r*² = 0.9924), indicating the operation of compensation effect in this reaction [14]. The value of isokinetic temperature evaluated [15, 16] from this plot is 1234±77 K. The correlation was tested and found genuine by Exner's criterion [17]. The isokinetic temperature, calculated from Exner's plot of log *k*₂ at 288 K versus log *k*₂ at 318 K (*r*² = 0.9998) is 1289±85 K. A linear isokinetic relationship is necessary condition for the validity of linear free energy relationship, which suggests that all the hydroxy acids are oxidized by the same mechanism.

Solvent effect: The rate constants of oxidation, *k*₂, in eighteen solvents (CS₂ was not considered, as the complete range of solvent parameters was not available) were correlated in terms of the linear solvation energy relationship (6) of Kamlet *et al.* [18].

$$\log k_2 = A_0 + p\pi^* + b\beta + a\alpha \quad (6)$$

In this equation π^* represents the solvent polarity, β the hydrogen bond acceptor basicities and α is the hydrogen bond donor acidity. A_0 is the intercept term. It may be mentioned here that out of the 18 solvents, 12 have a value of zero for α . The results of correlation analyses in terms of (6), a biparametric equation involving π^* and β , and separately with π^* and β are given below [(7 – 10)].

$$\log k_2 = -3.62 + (1.54 \pm 0.19)\pi^* + (0.09 \pm 0.16)\beta - (0.14 \pm 0.15)\alpha \quad (7)$$

$$r^2 = 0.8512; \text{ sd} = 0.18; n = 18; \Psi = 0.42$$

$$\log k_2 = -3.65 + (1.59 \pm 0.18)\pi^* + (0.04 \pm 0.15)\beta \quad (8)$$

$$r^2 = 0.8427; \text{ sd} = 0.18; n = 18; \Psi = 0.42$$

$$\log k_2 = -3.64 + (1.60 \pm 0.17)\pi^* \quad (9)$$

$$r^2 = 0.8419; \text{ sd} = 0.17; n = 18; \Psi = 0.41$$

$$\log k_2 = -2.72 + (0.33 \pm 0.35)\beta \quad (10)$$

$$r^2 = 0.0517; \text{ sd} = 0.42; n = 18; \Psi = 1.00$$

Here n is the number of data points and Ψ is the Exner's statistical parameter [19].

Kamlet's [18] triparametric equation explains *ca.* 85 % of the effect of solvent on the oxidation. However, by Exner's criterion (Fig. 3, Table 6) the correlation is not even satisfactory (*cf.* 5). The major contribution is of solvent polarity. It alone accounted for *ca.* 84 % of the data. Both β and α play relatively minor roles. The data on the solvent effect were analyzed in terms of Swain's equation [20] of cation- and anion-solvating concept of the solvents also (11).

$$\log k_2 = aA + bB + C \quad (11)$$




Figure 3. Exner's Isokinetic Relationship in the oxidation of Mandelic Acid by IDC.

Here A and B represent the anion-solvating and cation-solvating power. C is the intercept term. $(A + B)$ is postulated to represent the solvent polarity. The rates in different solvents were analyzed in terms of (12), separately with A and B and with $(A + B)$.

Table 6. Temperature dependence of the reaction constant

Temp./ K	$-\rho^*$	r^2	Sd	ψ
288	2.25±0.01	0.9999	0.006	0.01
298	2.16±0.02	0.9998	0.005	0.02
308	2.07±0.01	0.9989	0.009	0.03
318	1.98±0.02	0.9999	0.005	0.01

$$\log k_2 = 0.55 (\pm 0.05) A + 1.63 (\pm 0.04) B - 3.84 \quad (12)$$

$$r^2 = 0.9927; \text{sd} = 0.04; n = 19; \Psi = 0.09$$

$$\log k_2 = 0.32 (\pm 0.54) A - 2.72 \quad (13)$$

$$r^2 = 0.0205; \text{sd} = 0.44; n = 19; \Psi = 1.02$$

$$\log k_2 = 1.59 (\pm 0.10) B - 3.66 \quad (14)$$

$$r^2 = 0.9326; \text{sd} = 0.11; n = 19; \Psi = 0.27$$

$$\log k_2 = 1.27 \pm 0.14 (A + B) - 3.80 \quad (15)$$

$$r^2 = 0.8283; \text{sd} = 0.18; n = 19; \Psi = 0.43$$

The rates of oxidation of *p*-methylmandelic acid in different solvents showed an excellent correlation in Swain's equation (cf. 12) with the cation-solvating power playing the major role. In fact, the cation-solvation alone accounts for *ca* 99% of the data. The correlation with the anion-solvating power was very poor. The solvent polarity, represented by (A + B), also accounted for *ca.* 83 % of the data. In view of the fact that solvent polarity is able to account for *ca.* 83 % of the data, an attempt was made to correlate the rate with the relative permittivity of the solvent. However, a plot of $\log k_2$ against the inverse of the relative permittivity is not linear ($r^2 = 0.4773$; Sd = 0.32; $\Psi = 0.74$).

Correlation analysis of reactivity: The reaction rates of hydroxy acids were correlated with Hammett's σ values but did not yield very significant correlation. Though the rates of oxidation of substituted mandelic acids correlated well with Brown's σ^+ values [21], the reaction constant being negative (Table 5). The large negative reaction constants and correlation with σ^+ values indicate a carbo-cationic reaction centre in the transition state.


Mechanism: Absence of any effect of added acrylonitrile on the reaction discounts the possibility of a one-electron oxidation, leading to the formation of free radicals. The presence of a substantial kinetic isotope effect in the oxidation of mandelic acid confirms the cleavage of the α -C-H bond in the rate-determining step. The large negative reaction constant (-1.88 at 298 K) together with the excellent correlation with Brown's σ^+ values [21], point to a highly electron-deficient carbon centre in the transition state. The transition state thus approaches a carbocation in character. This is supported by the solvent effect also. Greater role played by the cation-solvating power of the solvents supported the postulation of a carbocationic transition state. Therefore, the correlation analysis of substituent and solvent effects on the oxidation of mandelic acid supports the mechanism involving a hydride-ion transfer via a chromate ester.

Kwart and Nickle [22] have shown that a study of the dependence of the kinetic isotope effect on temperature can be gainfully employed to resolve this problem. The data for protio and deuteriomandelic acids, fitted to the familiar expression $k_H/k_D = A_H/A_D \exp(E_a/RT)$ [23, 24] show a direct correspondence with the properties of a symmetrical transition state in which the activation energy difference (E_a) for k_H/k_D is equal to the zero-point energy difference for the respective C-H and C-D bonds ($\approx 4.5 \text{ kJ mol}^{-1}$) and the frequency factors and the entropies of activation of the respective reactions are nearly equal.


Bordwell [25] has documented very cogent evidence against the occurrence of concerted one-step biomolecular processes by hydrogen transfer and it is evident that in the present studies also the hydrogen transfer does not occur by an acyclic biomolecular process. It is well established that intrinsically concerted sigmatropic reactions, characterized by transfer of hydrogen in a cyclic transition state, are the only truly symmetrical processes involving a linear hydrogen transfer [26]. Littler [27] has also shown that a cyclic hydride transfer, in the oxidation of alcohols by Cr(VI), involves six electrons and, being a Huckel-type system, is an allowed process. Thus, the overall

mechanism is proposed to involve the formation of a chromate ester in a fast pre-equilibrium and then a decomposition of the ester in a subsequent slow step via a cyclic concerted symmetrical transition state leading to the product (Scheme 1 and 2). The observed negative value of entropy of activation also supports a polar transition state.

The observed negative entropy of activation also supports it. As the charge separation takes place, the charged ends become highly solvated. This results in an immobilization of a large number of solvent molecules, reflected in the loss of entropy [28].



Scheme 1



Scheme 2

APPLICATION

Oxidation reactions form an important part of organic chemistry. Kinetic studies of oxidation reactions of organic compounds have a bearing on the processes of metabolism also. It tells us about the rate enhancement or decrease due to field, resonance or steric factors of the substituents.

CONCLUSION

The reaction is proposed to proceed through a hydride-ion transfer from hydroxy acid to the oxidant. The hydride ion transfer mechanism is also supported by major role of cation-solvating power of the solvents. Both deprotonated and protonated forms of Imidazolium Dichromate are the reactive oxidizing species. Cleavage of α -C-H in the rate-determining step is taking place.

ACKNOWLEDGEMENTS

Thanks are due to the DST, New Delhi, India for financial support in the form of INSPIRE fellowship (to Ms. Gifty Sharma) and to Dr. Laszlo Kotai for re-synthesizing and characterization of IDC.

REFERENCES

- [1]. G. Cainelli and G. Cardillo, *Chromium oxidations in organic chemistry*, (Springer-Verlag, Berlin) Vol. 19 (1984).
- [2]. H. Firouzabadi, A. Sharifi, *Synthesis*, **1992**, 999.
- [3]. M. Li, M.E. Johnson, *Synth Commun*, **1995**, 25, 533.
- [4]. M.K. Mahanti, K.K. Banerji, *J. Indian Chem. Soc.*, **2002**, 79, 31.
- [5]. S. Kim, DC Lhim, *Bull. Chem. Soc. Jpn.*, **1986**, 59, 3297.
- [6]. A. Sharma, A. Meena J. Khatri, P. Swami, Vinita Sharma, *J. Applicable Chem.*, **2012**, 1(1), 78.
- [7]. L. Mathur, V. Ranga, G. Goswami, S. Agarwal, P.K. Sharma, *Eur. Chem. Bull.*, **2014**, 3(8) 828.
- [8]. A. Choudahry, S. Saraf, S. Puniya, Vinita Sharma, *J. Applicable Chem.*, **2016**, 5(4), 870.
- [9]. U. Songara, R. Kumar, Bhagyalaxmi, K. Vadera, P. K. Sharma, *J. Applicable Chem.*, **2016**, 5(5), 1225.
- [10]. K. K. Rathore, R. Sharma, J. Solanki, G. Sharma and Vinita Sharma, *J. Applicable Chem.*, **2018**, 7(3), 648.
- [11]. A. I. Vogel, *A text book of practical organic chemistry*, (Longmans, London) **1956**, 776.
- [12]. T. J. Kemp, W.A. Waters, *J. Chem. Soc.*, **1964**, 1192.
- [13]. D. D. Perrin, W. L. Armarego, D. R. Perrin, *Purification of organic Compounds*, Pergamon Press, Oxford, **1966**.
- [14]. L. Liu and W.E. Guo, *Chem. Review*, **2001**, 101, 673.
- [15]. J. E. Leffler, *J. Org. Chem.*, **1955**, 20, 1202; *J. Phys. Chem.*, **1964**, 29, 2199.
- [16]. R. C. Patterson, *J. Org. Chem.*, **1964**, 29, 3133.
- [17]. O. Exner, *Collect. Chem. Czech. Commun.*, **1977**, 38, 411.
- [18]. M. J. Kamlet, J. L. M. Abboud, M. H. Abraham, R. W. Taft, *J. Org. Chem.*, **1983**, 48, 2877.
- [19]. O. Exner, *Collect. Chem. Czech. Commun.*, **1966**, 31, 3222.
- [20]. C. G. Swain, S.H. Unger, N.R. Rosenquest, M.S. Swain, *J. Am. Chem. Soc.*, 1983, **105**, 492.
- [21]. H. C. Brown and Y. Okamoto, *J. Am. Chem. Soc.*, **1958**, 80, 4079.
- [22]. H. Kwart and J.H. Nickel, *J. Am. Chem. Soc.*, **1953**, 95, 3394.
- [23]. H. Kwart and M.C. Latimer, *J. Am. Chem. Soc.*, **1971**, 93, 3770.
- [24]. H. Kwart and J. Slutsky, *J. Chem. Soc. Chem. Commun*, **1972**, 1182.
- [25]. F. G. Bordwell, *Acc. Chem. Res.*, **1974**, 5, 374.
- [26]. R.W. Woodward and R. Hoffmann, *Angew. Chem. Int. Ed. Eng.*, **1969**, 8, 781.
- [27]. J. S. Littler, *Tetrahedron*, **1971**, 27, 81.
- [28]. E. S. Gould, *Mechanism and structure in organic chemistry*, (Holt, Rinehart & Winston, Inc. NY), **1964**.